

Zeta functions of finite Schreier graphs and their zig zag products

Asif Shaikh, Hemant Bhate

Abstract

We investigate the Ihara zeta functions of finite Schreier graphs Γ_n of the Basilica group. We show that Γ_{1+n} is 2 sheeted unramified normal covering of Γ_n , $\forall n \geq 1$ with Galois group $\frac{\mathbb{Z}}{2\mathbb{Z}}$. In fact, for any $n > 1, r \geq 1$ the graph Γ_{n+r} is 2^n sheeted unramified, non normal covering of Γ_r . In order to do this we give the definition of the *generalized replacement product* of Schreier graphs. We also show the corresponding results in zig zag product of Schreier graphs Γ_n with a 4 cycle.

Mathematics Subject Classification (2010): 05C31, 05C50, 05C76, 20E08

Keywords: Ihara zeta functions, Schreier graphs, zig zag product of graphs, Basilica groups.

1 INTRODUCTION

The main aim of this paper is to study the Ihara zeta functions of finite Schreier graphs Γ_n of the Basilica group as well as Ihara zeta functions of zig zag products of these Schreier graphs with a 4 cycle. The Ihara zeta function is a graph theoretic analogue of the Riemann zeta function $\zeta(s)$. It is defined for a connected graph $G = (V, E)$, for $t \in \mathbb{C}$, with $|t|$ sufficiently small, by

$$\zeta_G(t) = \prod_{[C] \text{ prime cycle in } G} (1 - t^{\nu(C)})^{-1} \quad (1)$$

where a prime $[C]$ in G is an equivalence class of tailless, backtrackless primitive cycles C in G , length of C is $\nu(C)$. See [8]. The connection of $\zeta_G(t)$ with the adjacency matrix A of G is given by Ihara-Bass's Theorem which says

$$\zeta_G(t)^{-1} = (1 - t^2)^{r-1} \det(I - At + Qt^2), \quad (2)$$

where $r = |E| - |V| - 1 = \text{rank of fundamental group of } G$ and Q is the diagonal matrix whose j^{th} diagonal entry is $Q_{jj} = d(v_j) - 1$, where $d(v_j)$ is the degree of the j^{th} vertex of G . The proof of the Ihara-Bass's theorem can be found in [8]. We say that finite graph $\tilde{G}$ is an unramified covering of a finite graph G if there is a covering map $\pi : \tilde{G} \rightarrow G$ which is an onto map such that for every vertex $u \in G$ and for every $v \in \pi^{-1}(u)$, the set of points adjacent to v in $\tilde{G}$ is mapped by π one-to-one, onto the vertices in G which are adjacent to u . A d -sheeted covering is a normal or Galois covering iff there are d graph automorphisms $\sigma : \tilde{G} \rightarrow \tilde{G}$ such that $\pi(\sigma(v)) = \pi(v)$, $\forall v \in \tilde{G}$. These automorphisms form the Galois group $\mathbb{G} = \text{Gal}(\tilde{G}|G)$. See [4, 8] for more details.

Our motivation comes from a question raised by A. Terras [8] of how zeta functions behave with respect to graph products, in particular, zig zag products. We have been able to prove the following:

If $n > 1$, then $\Gamma_{n+r} \mathbb{Z} C_4$ is the non normal covering of the graph $\Gamma_r \mathbb{Z} C_4$. In fact if $n = 1$ then $\Gamma_{1+r} \mathbb{Z} C_4$ is 2 sheeted normal covering of the graph $\Gamma_r \mathbb{Z} C_4$. Thus $\zeta_{(\Gamma_r \mathbb{Z} C_4)}(u)^{-1}$ divides $\zeta_{(\Gamma_{n+r} \mathbb{Z} C_4)}(u)^{-1}$, $\forall n$.

In order to prove the above result a new *generalized replacement product* $\Gamma_n \mathbb{G} \Gamma_r$ has been defined for two Schreier graphs Γ_n, Γ_r which gives the resultant graph $\Gamma_n \mathbb{G} \Gamma_r$ is also the Schreier graph Γ_{n+r} . Also,

If $n > 1$, then Γ_{n+r} is non normal covering of the graph Γ_r . In fact Γ_{1+r} is 2 sheeted normal covering of the graph Γ_r and $\zeta_{\Gamma_r}(u)^{-1}$ divides $\zeta_{\Gamma_{n+r}}(u)^{-1}$, $\forall n$.

This paper is organized as follows: In section 2, we define Schreier graphs of the Basilica group and give several examples. This section also contains the definition of *generalized replacement product* and the computations of Ihara zeta functions of these Schreier graphs using Artin L functions. This section ends with some results which will be needed in the next section. Section 3 starts with the definition of zig zag product of two graphs see [7, 3]. Computations of Ihara zeta functions of zig zag product of Schreier graphs with a 4 cycle are also presented. We also prove the corresponding results in zig zag product of graphs.

2 SCHREIER GRAPHS OF THE BASILICA GROUP

Let $X = \{0, 1\}$ be a binary alphabet. Denote X^0 the set consisting of the empty word and by $X^n = \{w = x_1 x_2 \cdots x_n : x_i \in X\}$ the set of words of length n over the alphabet X , for each $n \geq 1$. The Basilica group B acting on X^n is the group generated by three state automaton. The states a and b of the automaton are the generators of the group. The action of a and b is given by

$$a(0w) = 0b(w), a(1w) = 1w, b(0w) = 1a(w), b(1w) = 0w, \quad \forall w \in X^n.$$

For each $n \geq 1$, let Γ_n be the *Schreier graph* associated with the action of B on X^n with $V(\Gamma_n) = X^n$ and two vertices v, v' are connected by an edge labeled by s near v and by s^{-1} near v' if $s(v) = v'$ (i.e. $s^{-1}(v') = v$), where $s \in \{a^{\pm 1}, b^{\pm 1}\}$. As the action of B on X^n is transitive, the graph Γ_n is a 4 regular connected graph on 2^n vertices and labels near v are given by $a^{\pm 1}, b^{\pm 1}$ for every $v \in X^n$. We denote $s(v)$ by v^s , where $v \in X^n, s \in \{a^{\pm 1}, b^{\pm 1}\}$.

More information about such finite and infinite Schreier graphs of Basilica group




Figure 1: The graph $\Gamma_1, \Gamma_2, \Gamma_3$ and Γ_4 are the Schreier graphs of the Basilica group over X, X^2, X^3 and X^4 respectively.

can be found in [2, 3]. The complete classification (up to isomorphism) of the

limiting case of infinite Schreier graphs associated with the Basilica group acting on the binary tree, in terms of the infinite binary sequence is given by D. D'Angeli, A. Donno, M. Matter and T. Nognibeda[2]. D. D'Angeli, A. Donno and E. Sava-Huss[3] have used Schreier graphs of Basilica groups in computations of zig zag product of graphs. The Basilica group belongs to the important class of self-similar groups and was introduced by R. Grigorchuk and A. Żuk[5].

Example 2.1. *The first four finite Schreier graphs of Basilica group are as in Figure 1.*

The substitution rules [3, 6] can be used to construct the Schreier graph Γ_{n+1} from the graph Γ_n . In the next sub section we give the new definition of the product of two Schreier graphs, we call it as *generalized replacement product* of Schreier graphs.

2.1 Generalized replacement product

The replacement product of two graphs is well known in literature. If G_1 and G_2 are two regular graphs with the regularity d_1 and d_2 respectively, then their replacement product $G_1 \circledast G_2$ is again a regular graph with regularity $d_2 + 1$. The details about this product can be found in [3, 1]. Below is the generalization of this product, in which both G_1 and G_2 along with the resultant graph have same regularity. The motivation of generalized version is the following: Using any two Schreier graphs of the Basilica group we should be able to produce the resultant graph to be Schreier graph of the Basilica group.

Let Γ_n and Γ_r be two Schreier graphs of Basilica group. To define their generalized replacement product, we first choose the spanning subgraph Γ'_r of the graph Γ_r such that $E(\Gamma'_r) = E(\Gamma_r) \setminus \{e_a, e_b\}$ where e_a and e_b are the edges of the vertex $u_0 = 0^r \in \Gamma_r$ with labels a and b near u_0 respectively. Note that the vertex $u_0 = 0^r$ we mean the word of length r and having all alphabets are 0. Let $a(u_0) = u_0^a$ and $b(u_0) = u_0^b$ be the vertices of Γ'_r which are adjacent to u_0 in Γ_r by the edges e_a and e_b respectively.

Thus every vertex from the set $X^r \setminus \{u_0, u_0^a, u_0^b\}$ has degree $d = 4$ and

$$d(u_0) = 2, d(u_0^a) = d(u_0^b) = 3$$

$$(4 - d(u_0)) + (4 - d(u_0^a)) + (4 - d(u_0^b)) = 2 + 1 + 1 = 4$$

Note that in above equation sum of LHS should be the regularity d of the graph.

Notice also that in the Schreier graphs Γ_n of the Basilica group, if $e = \{v, v'\}$

is an edge which has color say s near v and s^{-1} near v' , then the *rotation map* $\text{Rot} : X^n \times D \rightarrow X^n \times D$ is defined by

$$\text{Rot}_{\Gamma_n}(v, s) = (v', s^{-1}), \quad \forall v, v' \in X^n, \quad s, s^{-1} \in D = \{a^{\pm 1}, b^{\pm 1}\}.$$

Definition 2.1. *The generalized replacement product $\Gamma_n \textcircled{g} \Gamma_r$ is the 4 regular graph with vertex set $X^{n+r} = X^n \times X^r$, and whose edges are described by the following rotation map: $\forall v \in X^n$*

$$\text{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u_0), a) = \begin{cases} ((v^a, u_0^a), a^{-1}) & \text{if } r \text{ is even} \\ ((v^b, u_0^a), a^{-1}) & \text{if } r \text{ is odd} \end{cases} \quad (3)$$

$$\text{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u_0), b) = \begin{cases} ((v^b, u_0^b), b^{-1}) & \text{if } r \text{ is even} \\ ((v^a, u_0^b), b^{-1}) & \text{if } r \text{ is odd} \end{cases} \quad (4)$$

$$\text{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), s) = ((v, u^s), s^{-1}), \quad \forall u \in X^r \setminus \{u_0\} \quad s \in \{a^{\pm 1}, b^{\pm 1}\} \quad (5)$$

We call edges given in (3) as lifts of the edge e_a and in (4) as lifts of the edge e_b .

One can imagine that the vertex set X^{n+r} of the graph $\Gamma_n \textcircled{g} \Gamma_r$ is partitioned into the sheets, which are indexed by the vertices of Γ_n , where by definition of the v^{th} sheet, for $v \in X^n$, consists of the vertices $\{(v, u) | u \in X^r\}$. Within this construction the idea is to put the copy of Γ'_r around each vertex v of Γ_n , while keeping all the edges of Γ'_r as it is and the Γ_n can be thought of using the above given rotations in the equations (3) and (4). The connectedness of Γ'_r and Γ_n guarantees about the connectedness of the graph $\Gamma_n \textcircled{g} \Gamma_r$.

If an edge $e = \{v, v'\}$ in Γ_n has colors s and s^{-1} near v and v' respectively, then we call the edge e as s -edge. Hence we can say that in the graph Γ_n , there are two types of edges i.e. a -edges and b -edges.

Remark 2.1. *In the Figure 2,3 and 4 one can observe that the number of edges in the graph Γ_n and the number of edges which are lifts of e_a and e_b in $\Gamma_n \textcircled{g} \Gamma_r$ are the same. If we imagine that the whole sheet as a single vertex along with the lift edges interestingly we get the graph Γ_n . Hence we say the following:*

1. *If r is even then the edges which are lifts of e_a and e_b in $\Gamma_n \textcircled{g} \Gamma_r$ are a and b edges in Γ_n respectively.*
2. *If r is odd then the edges which are lifts of e_a and e_b in $\Gamma_n \textcircled{g} \Gamma_r$ are b and a edges in Γ_n respectively.*
3. *The edges which are not lifts of e_a and e_b in $\Gamma_n \textcircled{g} \Gamma_r$ are edges in Γ_r , so we call them sheet edges.*




Figure 2: We choose $u_0 = 110$. So the graph $\Gamma_1 \otimes \Gamma_3$ is still normal 2 sheeted cyclic covering of the graph Γ_3 but it is not isomorphic to Γ_4 . The 2 sheets of Γ_3 are copies of the spanning subgraph Γ'_3 (dashed lines). Black continuous lines/loops in $\Gamma_1 \otimes \Gamma_3$ are lifts of e_a or e_b .

Note that above definition of $\Gamma_n \otimes \Gamma_r$ depends on the spanning subgraph of the second graph. If a different choice of spanning subgraph is taken, then the graph $\Gamma_n \otimes \Gamma_r$ need not be Schreier graph of the Basilica group.

Example 2.2. If we choose a vertex of Γ_r other than 0^r as u_0 and then corresponding spanning subgraph Γ'_r with edge set, $E(\Gamma_r) \setminus \{e_a, e_b\}$ where e_a and e_b are the edges of the vertex u_0 with labels a and b respectively near u_0 , then the graph $\Gamma_n \otimes \Gamma_r$ need not be Schreier graph of the Basilica group. See Figure 2.

The following result will show that the graph $\Gamma_n \otimes \Gamma_r$ is actually a Schreier graph under the action of Basilica group on the set X^{n+r} .

Proposition 2.1. Let $n, r \geq 1$, then the following holds:

1. The graph $\Gamma_{n+r} \simeq \Gamma_n \otimes \Gamma_r \simeq \Gamma_r \otimes \Gamma_n$ is a Schreier graph under the action of Basilica group on the set X^{n+r} .
2. Γ_{n+r} is an unramified, 2^n sheeted graph covering of Γ_r .
3. $\Gamma_n \otimes \Gamma_r$ is non normal covering of the graph Γ_r if $n > 1$.
4. Γ_{1+r} is 2 sheeted normal covering of the graph Γ_r .

Proof. Let $n, r \geq 1$.

1. Let us define the map $f : \Gamma_n \otimes \Gamma_r \rightarrow \Gamma_{n+r}$ by $f(v, u) = uv$, where $v \in X^n$ and $u \in X^r$. To show f is adjacency preserving map, notice that there are two types of edges in $\Gamma_n \otimes \Gamma_r$, first type which contains those edges which are lifts of e_a or e_b , i.e. edges which are described in the equation (3) or (4) and second type contains edges which are not lifts of e_a or e_b , i.e. edges which are described in the equation (5).

Edges of the first type:

Note that lift of e_a connect (v, u_0) to (v', u_0^a) and lift of e_b connect (v, u_0) to (v'', u_0^b) where

$$v' = \begin{cases} v^a & \text{if } r \text{ is even} \\ v^b & \text{if } r \text{ is odd} \end{cases}$$

$$v'' = \begin{cases} v^b & \text{if } r \text{ is even} \\ v^a & \text{if } r \text{ is odd} \end{cases}$$

Let given edge be a lift of e_a . In other words, let (v, u_0) be adjacent to (v', u_0^a) in $\Gamma_n \otimes \Gamma_r$, i.e.

$$\Leftrightarrow \mathbf{Rot}_{\Gamma_n \otimes \Gamma_r}((v, u_0), a) = ((v', u_0^a), \alpha^{-1})$$

Then by action of the Basilica group B on X^{n+r} ,

$$\Leftrightarrow a(u_0 v) = \begin{cases} u_0^a v^a & \text{if } r \text{ is even} \\ u_0^a v^b & \text{if } r \text{ is odd} \end{cases}$$

$$\Leftrightarrow a(u_0 v) = u_0^a v', v' = \begin{cases} v^a & \text{if } r \text{ is even} \\ v^b & \text{if } r \text{ is odd} \end{cases}$$

$$\Leftrightarrow \mathbf{Rot}_{\Gamma_{n+r}}(u_0 v, a) = (u_0^a v'), v' = \begin{cases} v^a & \text{if } r \text{ is even} \\ v^b & \text{if } r \text{ is odd} \end{cases}$$

$\Leftrightarrow u_0 v$ is adjacent to $u_0^a v'$ in Γ_{n+r} which shows that f is adjacency preserving map. The argument is similar if the given edge is a lift of e_b .

Edges of the second type:

The edges of $\Gamma_n \otimes \Gamma_r$ which are not lifts of e_a or e_b connect (v, u) to (v, u^s) where $u, u^s \neq u_0$. Suppose given edge is of this type.

$$\Leftrightarrow \mathbf{Rot}_{\Gamma_n \otimes \Gamma_r}((v, u), s) = ((v, u^s), s^{-1}), s \in \{a^{\pm 1}, b^{\pm 1}\}, u, u^s \neq u_0.$$

As $u \neq u_0 = 0^r$ means u has at least one “1” as an alphabet, therefore by definition of Basilica group elements we have

$$s(uv) = s(u)v = u^s v, s \in \{a^{\pm 1}, b^{\pm 1}\}$$

Hence uv is adjacent to $u^s v$ in Γ_{n+r} which gives f is adjacency preserving map. By definition of f it is not difficult to that f is bijection map. Therefore f^{-1} exists and by reverse implication it shows that f^{-1} is also adjacency preserving map. As f bijection so as f^{-1} . Thus f and f^{-1} both are isomorphisms.

2. By definition 2.1, $\Gamma_{n+r} \simeq \Gamma_n \mathbin{\text{\textcircled{G}}} \Gamma_r$ contains 2^n sheets of the graph Γ_r . To show that Γ_{n+r} is an unramified cover of the graph Γ_r , we define the map $\pi : \Gamma_{n+r} \rightarrow \Gamma_r$ by $\pi(uv) = u$. We now show that π is a covering map i.e. π is sending neighborhoods of Γ_{n+r} one-to-one and onto neighborhoods of Γ_r .

Suppose $u_0 v$ is adjacent to $u_0^s v'$ in Γ_{n+r} where $s \in \{a, b\}$.

By definition of π , $\pi(u_0 v) = u_0$ and $\pi(u_0^s v') = u_0^s$ but $u_0^s = s(u_0)$ which means u_0 is adjacent to $s(u_0) = u_0^s$ in Γ_r so $\pi(u_0 v)$ is adjacent to $\pi(u_0^s v')$ in Γ_r .

Let

$$\mathbf{N}_{\Gamma_{n+r}}(u_0 v) = \{u_0^a v', u_0^b v'', u_0^{a^{-1}} v, u_0^{b^{-1}} v\},$$

then

$$\begin{aligned} \pi(\mathbf{N}_{\Gamma_{n+r}}(u_0 v)) &= \pi(\{u_0^a v', u_0^b v'', u_0^{a^{-1}} v, u_0^{b^{-1}} v\}) \\ &= \{u_0^a, u_0^b, u_0^{a^{-1}}, u_0^{b^{-1}}\} = \mathbf{N}_{\Gamma_r}(u_0) \end{aligned}$$

Now suppose uv is adjacent to $u^s v$ in Γ_{n+r} where $s \in \{a^{\pm 1}, b^{\pm 1}\}$ and $u, u^s \neq u_0$.

By definition of π , $\pi(uv) = u$ and $\pi(u^s v) = u^s$ but $u^s = s(u)$ in Γ_r gives u is adjacent to u^s in Γ_r .

Let

$$\mathbf{N}_{\Gamma_{n+r}}(uv) = \{u^a v, u^b v, u^{a^{-1}} v, u^{b^{-1}} v\},$$

then

$$\begin{aligned} \pi(\mathbf{N}_{\Gamma_{n+r}}(uv)) &= \pi(\{u^a v, u^b v, u^{a^{-1}} v, u^{b^{-1}} v\}) \\ &= \{u^a, u^b, u^{a^{-1}}, u^{b^{-1}}\} = \mathbf{N}_{\Gamma_r}(u) \end{aligned}$$

$\implies \pi$ is a covering map. Thus Γ_{n+r} is an unramified covering of Γ_r .

3. (Proof by contradiction) Assume that Γ_{n+r} is normal covering of Γ_r , where $n > 1$. The Galois group $\mathbb{G} = \text{Gal}(\Gamma_{n+r} | \Gamma_r) = \text{Gal}(\Gamma_n \mathbin{\text{\textcircled{G}}} \Gamma_r | \Gamma_r)$ is generated by permutations corresponding to the Frobenius automorphism σ at e_a and e_b . See [8]

i.e.

$$\mathbb{G} = \langle \sigma(e_a), \sigma(e_b) \rangle$$

The permutations $\sigma(e_a)$ and $\sigma(e_b)$ arises under the action of Basilica group elements a and b on the (ordered) set X^n respectively when we consider

Γ_{n+r} as unramified covering of Γ_r .

As Γ_{n+r} has 2^n sheets of Γ_r , we indexed these sheets by the set $\{1, 2, \dots, 2^n\} \sim X^n$. If two sheets say i^{th} and j^{th} are connected by the lift of say e_a then we write $\sigma(e_a)(i) = j$ similarly we can write the permutation corresponds to $\sigma(e_b)$.

The permutations $\sigma(e_a), \sigma(e_b)$ can be written directly using action of Basilica group elements a and b on the set X^n respectively i.e. suppose $v_i, v_j \in X^n$, with $a(v_i) = v_j$ then we write $\tilde{\sigma}(e_a)(i) = j$. By Remark 2.1, $\sigma \equiv \tilde{\sigma}$. Therefore the product $\sigma(e_a) \cdot \sigma(e_b)$ comes from the action of ba on X^n . Using induction, it can be shown as in [3] that the action of ba on X^n has order 2^n . This shows that the Galois group $\mathbb{G}$ has an element $(\sigma(e_a) \cdot \sigma(e_b))$ which is different from the generators $\sigma(e_a)$ and $\sigma(e_b)$ and which has order 2^n . Hence Galois group $\mathbb{G}$ has order bigger than 2^n which is a contradiction.

4. Suppose $n = 1$, therefore Γ_{1+r} is 2 sheeted covering of Γ_r so we first show that $|\mathbb{G}| = 2$. Denote two sheets Γ'_r of Γ_{1+r} by 1 and 2.

Recall that the Galois group $\mathbb{G} = \text{Gal}(\Gamma_{1+r}|\Gamma_r) = \langle \sigma(e_a), \sigma(e_b) \rangle$. See [8]. By the definition of $\Gamma_{1+n} \simeq \Gamma_1 \otimes \Gamma_n$ we have the following:

If r is even then

$$\sigma(e_a) = (1), \sigma(e_b) = (1\ 2)$$

and if r is odd then

$$\sigma(e_a) = (1\ 2), \sigma(e_b) = (1)$$

Therefore

$$\mathbb{G} = \langle \sigma = (1\ 2) \rangle, \forall r.$$

To show that Γ_{1+r} is 2 sheeted normal covering of Γ_r it remains to show that $\pi \circ \sigma \equiv \pi$ where $\sigma \in \mathbb{G}$ is the non identity Frobenius automorphism with $\sigma^2 = id$. If u is any vertex of Γ_r so let $u0$ be any vertex of the first sheet and $u1$ be any vertex of the second sheet.

$$(\pi \circ \sigma)(u0) = \pi(\sigma(u0)) = \pi(u1) = u$$

$$(\pi \circ \sigma)(u1) = \pi(\sigma(u1)) = \pi(u0) = u,$$

and

$$\pi(u1) = u = \pi(u0),$$

Therefore

$$\pi \circ \sigma \equiv \pi.$$

□

Corollary 2.1. *The spectrum of Γ_r is contained in the spectrum of Γ_{1+r} . Moreover The zeta function $\zeta_{\Gamma_r}(t)^{-1}$ divides $\zeta_{\Gamma_{1+r}}(t)^{-1}$.*

Proof. This is direct consequence of the proof of the Proposition 13.10, p.no. 110 given in [8]. $\square$

Let $\tilde{Y}$ be the normal covering of Y . The Artin L function associated to the representation ρ of $\mathbb{G} = \text{Gal}(\tilde{Y}|Y)$ can be defined by a product over prime cycles in Y as

$$L(u, \rho, \tilde{Y}|Y) = \prod_{[c] \text{ prime in } Y} \det(I - \rho(\text{Frob}(\tilde{C}))) u^{\nu(C)}^{-1}$$

where $\tilde{C}$ denotes any lift of C to $\tilde{Y}$ and $\text{Frob}(\tilde{C})$ denotes the Frobenius automorphism defined by

$$\text{Frob}(\tilde{C}) = ji^{-1},$$

if $\tilde{C}$ starts on $\tilde{Y}$ -sheet labeled by $i \in \mathbb{G}$ and ends on $\tilde{Y}$ -sheet labeled by $j \in \mathbb{G}$. As in Proposition 2.1 of [4], the adjacency matrix of $\tilde{Y}$ can be block diagonalized where the blocks are of the form

$$A_\rho = \sum_{g \in \mathbb{G}} A(g) \otimes \rho(g), \quad (6)$$

each taken $d_\rho = \text{degree of } \rho \text{ times}$, $\rho \in \hat{\mathbb{G}}$ irreducible representation of $\mathbb{G}$ and $A(g)$ is defined in following formula:

Suppose Y has m vertices. Define the $m \times m$ matrix $A(g)$ for $g \in \mathbb{G}$ by defining i, j entry to be

$$A(g)_{i,j} = \text{the number of edges in } \tilde{Y} \text{ between } (1, e) \text{ to } (j, g), \quad (7)$$

where e denotes the identity in $\mathbb{G}$.

By setting $Q_\rho = Q \otimes I_{d_\rho}$, with $d_\rho = \deg \rho$, we have the following analogue of formula (2):

$$L(t, \rho, \tilde{Y}|Y)^{-1} = (1 - t^2)^{(r-1)d_\rho} \det(I - A_\rho t + Q_\rho t^2). \quad (8)$$

See [8].

Thus we have zeta functions of $\tilde{Y}$ factors as follows

$$\zeta_{\tilde{Y}}(t) = \prod_{\rho \in \hat{\mathbb{G}}} L(t, \rho, \tilde{Y}|Y)^{d_\rho} \quad (9)$$

See Corollary 18.11 of [8].

The matrices A_1 and A_σ are called as the Artinized adjacency matrices.

We label the vertices using the Table 1.




Figure 3: The graph $\Gamma_1 \otimes \Gamma_2$ is a normal 2 sheeted cyclic covering of the graph Γ_2 . The 2 sheets of Γ_2 are copies of the spanning subgraph Γ'_2 (dashed lines). The map f is a graph isomorphism between $\Gamma_1 \otimes \Gamma_2$ and Γ_3 .

Table 1: Vertex labeling for Γ_2

label	1	2	3	4
vertex	11	01	00	10

Example 2.3. The covering $\Gamma_3 \simeq \Gamma_1 \circledast \Gamma_2$ over Γ_2 given in the Figure 3 is a normal covering. We obtain spanning subgraph of the $Y = \Gamma_2$ by cutting edges $e_a = \{3, 2\}, e_b = \{3, 4\}$. This gives dashed line spanning subgraph Γ'_2 of the Γ_2 . So we draw the covering graph $\tilde{Y} = \Gamma_3 \simeq \Gamma_1 \circledast \Gamma_2$ by placing 2 sheets of spanning subgraph Γ'_2 of $Y = \Gamma_2$ and labeling each sheet given in Table 2. Connections between 2 sheets in the cover graph $\tilde{Y} = \Gamma_3$ are given in the Table 3. In this case the representations of cyclic Galois group $\mathbb{G} = \{1, \sigma\}$ are the trivial representation $\rho_0 = 1$ and the representation ρ defined by $\rho(1) = 1, \rho(\sigma) = -1$. So $Q_\rho = 2I_4$. There are two cases.

Table 2: Notations for sheets

Vertex set of $\Gamma_1 \circledast \Gamma_2$	Vertex set of Γ_3	Group element
$\{1, 2, 3, 4\} \sim \{(0, 11), (0, 01), (0, 00), (0, 10)\}$	$\{110, 010, 000, 100\}$	$1 = id$
$\{1', 2', 3', 4'\} \sim \{(1, 11), (1, 01), (1, 00), (1, 10)\}$	$\{111, 011, 001, 101\}$	$\sigma \neq id$

Table 3: Connections between sheet 1 and 2 in $\Gamma_3 \simeq \Gamma_1 \circledast \Gamma_2$

Vertex	adjacent vertices in $\Gamma_3 \simeq \Gamma_1 \circledast \Gamma_2$
1	1, 2
2	1, 3
3	2, 4, 4'
4	3, 4, 3'

Case 1 The trivial representation $\rho_0 = 1$

Here $A_1 = A(1) + A(\sigma) = A_Y$ where

$$A(1) = \begin{pmatrix} 2 & 2 & 0 & 0 \\ 2 & 0 & 2 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix}, A(\sigma) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

and A_1 is the adjacency matrix A_Y .

Case 2 The representation ρ

$$A_\sigma = A(1) - A(\sigma) = \begin{pmatrix} 2 & 2 & 0 & 0 \\ 2 & 0 & 2 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}$$

Reciprocals of L functions for $\Gamma_3|\Gamma_2$.

1) For A_1

$$\zeta_{\Gamma_2}(t)^{-1} = L(t, A_1, \Gamma_3|\Gamma_2)^{-1} = (1-t^2)^4(t-1)(3t-1)(3t^2+1)(9t^4-2t^2+1).$$

2) For A_σ

$$L(t, A_\sigma, \tilde{Y}|Y)^{-1} = (1-t^2)^4(3t^2-2t+1)(27t^6-18t^5+3t^4-4t^3+t^2-2t+1)$$

By equation (9) we have

$$\begin{aligned} \zeta_{\Gamma_3}(t)^{-1} &= L(t, A_1, \tilde{Y}|Y)^{-1} L(t, A_\sigma, \tilde{Y}|Y)^{-1} \\ &= (1-t^2)^8(t-1)(3t-1)(3t^2+1)(3t^2-2t+1)(9t^4-2t^2+1) \\ &\quad (27t^6-18t^5+3t^4-4t^3+t^2-2t+1) \end{aligned}$$

Example 2.4. The coverings $\Gamma_5 \simeq \Gamma_3 \circledast \Gamma_2$ over Γ_2 given in the Figure 4 is non normal covering. We follow the labeling the vertices given in Table 1. We obtain spanning subgraph of the Γ_2 by cutting edges $e_a = \{3, 2\}, e_b = \{3, 4\}$. This gives dashed line spanning subgraph Γ'_2 of the Γ_2 . So we draw the covering graph $\Gamma_5 \simeq \Gamma_3 \circledast \Gamma_2$ by placing 8 sheets of spanning subgraph Γ'_2 of Γ_2 and labeling each with notations given in Table 4 and connections between first sheet with other sheets are given in Table 5.

Here we assume that $\Gamma_3 \circledast \Gamma_2$ is normal covering of Γ_2 .

$$\sigma(e_a) = (2\ 3)(6\ 7), \sigma(e_b) = (1\ 2)(3\ 5\ 6\ 4)(7\ 8)$$

and

$$\sigma(e_a) \circ \sigma(e_b) = (1\ 3\ 5\ 7\ 8\ 6\ 4\ 2)$$

Hence $\mathbb{G} = \text{Gal}(\Gamma_3 \circledast \Gamma_2|\Gamma_2) = \langle \sigma(e_a), \sigma(e_b) \rangle$ has an element of order 8. Thus

$$|\mathbb{G}| > 8$$

is a contradiction.




Figure 4: The graph $\Gamma_3 \otimes \Gamma_2$ is a non normal 8 sheeted covering of the graph Γ_2 . The 8 sheets of Γ_2 are copies of the spanning subgraph Γ'_2 (dashed lines). The graph $\Gamma_2 \otimes \Gamma_3$ is isomorphic to the graph Γ_5 .

Table 4: Notations for sheets

Vertex set of $\Gamma_3 \circledast \Gamma_2$	Sheet index
$\{1, 2, 3, 4\} \sim \{(110, 11), (110, 01), (110, 00), (110, 10)\}$	1^{st}
$\{1', 2', 3', 4'\} \sim \{(010, 11), (010, 01), (010, 00), (010, 10)\}$	2^{nd}
$\{1'', 2'', 3'', 4''\} \sim \{(000, 11), (000, 01), (000, 00), (000, 10)\}$	3^{rd}
$\{1''', 2''', 3''', 4'''\} \sim \{(100, 11), (100, 01), (100, 00), (100, 10)\}$	4^{th}
$\{1^{(5)}, 2^{(5)}, 3^{(5)}, 4^{(5)}\} \sim \{(101, 11), (101, 01), (101, 00), (101, 10)\}$	5^{th}
$\{1^{(6)}, 2^{(6)}, 3^{(6)}, 4^{(6)}\} \sim \{(001, 11), (001, 01), (001, 00), (001, 10)\}$	6^{th}
$\{1^{(7)}, 2^{(7)}, 3^{(7)}, 4^{(7)}\} \sim \{(011, 11), (011, 01), (011, 00), (011, 10)\}$	7^{th}
$\{1^{(8)}, 2^{(8)}, 3^{(8)}, 4^{(8)}\} \sim \{(111, 11), (111, 01), (111, 00), (111, 10)\}$	8^{th}

Table 5: Connections between sheet 1 and other sheets in $\Gamma_5 \simeq \Gamma_3 \circledast \Gamma_2$

Vertex	adjacent vertices in $\Gamma_5 \simeq \Gamma_3 \circledast \Gamma_2$
1	1, 2
2	1, 3
3	2, 4, 4'
4	3, 4, 3'

$\implies \Gamma_3 \circledast \Gamma_2$ is non normal covering of Γ_2 .

Proposition 2.2. Let $\Gamma_n \circledast \Gamma_r$ with $n > 1$ be the unramified, non normal covering of Γ_r then following holds:

1. If r is an even and $av = v$ in Γ_n then the alternate action of a and b^{-1} on $y_0 = (v, u_0)$ i.e.

$$\begin{aligned} y_{2p+1} &= ay_{2p}, \quad p = 0, 1, \dots, (2^r - 1) \\ y_{2p+2} &= b^{-1}y_{2p+1}, \quad p = 0, 1, \dots, (2^r - 2) \end{aligned} \quad (10)$$

produces spanning path on v^{th} sheet of $\Gamma_n \circledast \Gamma_r$ which visits every vertex of the v^{th} sheet twice.

2. If r is an odd and $av = v$ in Γ_n then the alternate action of b and a^{-1} on $y_0 = (v, u_0)$ i.e.

$$\begin{aligned} y_{2p+1} &= by_{2p}, \quad p = 0, 1, \dots, (2^r - 1) \\ y_{2p+2} &= a^{-1}y_{2p+1}, \quad p = 0, 1, \dots, (2^r - 2) \end{aligned} \quad (11)$$

produces spanning path on v^{th} sheet of $\Gamma_n \textcircled{g} \Gamma_r$ which visits every vertex of the v^{th} sheet twice.

3. Let r be any positive number with $av \neq v$ in Γ_n , then alternate actions of any of the above type goes outside the v^{th} sheet.

Proof. 1. As r is an even and $av = v$ in Γ_n , then by definition of generalized replacement product, v^{th} sheet in $\Gamma_n \textcircled{g} \Gamma_r$ contains lift of the edge e_a which connects (v, u_0) to (v, u_0^a) in $\Gamma_n \textcircled{g} \Gamma_r$. i.e. v^{th} sheet is actually a copy of the graph $\Gamma_r \setminus \{e_b\}$. So we can write

$$\mathbf{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), a) = ((v, u^a), a^{-1}), \forall u \in V_{\Gamma_r} \quad (12)$$

$$\mathbf{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), b^{-1}) = ((v, u^{b^{-1}}), b), \forall u \in V_{\Gamma_r} \setminus \{u_0^b\} \quad (13)$$

Therefore the simultaneous actions of a and b^{-1} as stated in (10), produces a path which visits the following vertices.

$$\begin{aligned} y_0 &= (v, u_0), \\ y_1 &= ay_0 = a(v, u_0) = (v, a(u_0)) = (v, u_1), \\ y_2 &= b^{-1}y_1 = b^{-1}(v, u_1) = (v, b^{-1}u_1) = (v, u_2), \\ &\dots y_{2r+1-1} = ay_{2r+1-2} = (v, u_{2r+1-1}) \end{aligned} \quad (14)$$

One can even prove that $(v, u_{2r+1-1}) = (v, u_0^b)$ i.e. $u_{2r+1-1} = u_0^b$. Using induction it has shown as in [3], that action of $b^{-1}a$ has order 2^r on $V(\Gamma_r) = X^r$. Thus

$$(b^{-1}a)^{2^r} u_0 = u_0$$

i.e.

$$\underbrace{b^{-1}ab^{-1}a \dots b^{-1}a}_{2^r \text{ times}} u_0 = u_0$$

But

$$u_{2r+1-1} = a(b^{-1}a)^{2^r-1} u_0 \implies b^{-1}(u_{2r+1-1}) = b^{-1}a(b^{-1}a)^{2^r-1} u_0 = u_0$$

This gives,

$$u_{2r+1-1} = u_0^b \implies (v, u_{2r+1-1}) = (v, u_0^b)$$

The right hand side of equation (14), shows that all y_i 's are vertices of v^{th} sheet of the graph $\Gamma_n \textcircled{g} \Gamma_r$ which are counted twice. Hence we get the spanning path on v^{th} sheet of $\Gamma_n \textcircled{g} \Gamma_r$ which visits every vertex of the v^{th} sheet twice.

2. As r is an odd number and $av = v$ in Γ_n , then by definition of generalized replacement product, v^{th} sheet in $\Gamma_n \textcircled{g} \Gamma_r$ contains lift of the edge e_b which connects (v, u_0) to (v, u_0^b) in $\Gamma_n \textcircled{g} \Gamma_r$. i.e. v^{th} sheet is actually a copy of the graph $\Gamma_r \setminus \{e_a\}$. So we can write

$$\mathbf{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), b) = ((v, u^b), b^{-1}), \forall u \in V(\Gamma_r) \quad (15)$$

$$\mathbf{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), a^{-1}) = ((v, u^{a^{-1}}), a), \forall u \in V(\Gamma_r) \setminus \{u_0^a\} \quad (16)$$

Therefore the simultaneous actions of b and a^{-1} as stated in (11), produces a path which visits the following vertices.

$$\begin{aligned} y_0 &= (v, u_0), \\ y_1 &= by_0 = b(v, u_0) = (v, b(u_0)) = (v, u_1), \\ y_2 &= a^{-1}y_1 = a^{-1}(v, u_1) = (v, a^{-1}u_1) = (v, u_2), \\ \dots y_{2r+1-1} &= by_{2r+1-2} = (v, u_{2r+1-1}) = (v, u_0^a) \end{aligned} \quad (17)$$

The right hand side of equation (17), shows that all y_i 's are vertices of v^{th} sheet of the graph $\Gamma_n \textcircled{g} \Gamma_r$ which are counted twice. Hence we get the spanning path on v^{th} sheet of $\Gamma_n \textcircled{g} \Gamma_r$ which visits every vertex of the v^{th} sheet twice.

3. Let $n > 1$ be positive integer, and $av \neq v \in \Gamma_n$. The lifts of e_a and e_b go from v^{th} sheet to another sheets. i.e. These lifts connect (v, u_0) to other sheets and hence the v^{th} sheet is actually a copy of the graph $\Gamma_r \setminus \{e_a, e_b\}$, therefore we have

$$\mathbf{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), a^{-1}) = ((v, u^{a^{-1}}), a), \forall u \in V(\Gamma_r) \setminus \{u_0^a\} \quad (18)$$

$$\mathbf{Rot}_{\Gamma_n \textcircled{g} \Gamma_r}((v, u), b^{-1}) = ((v, u^{b^{-1}}), b), \forall u \in V(\Gamma_r) \setminus \{u_0^b\} \quad (19)$$

If r is even, the action of a on (v, u_0) gives the vertex (v^a, u_0^a) but $v^a = a(v) \neq v \in \Gamma_n$. Therefore the action of a on (v, u_0) doesn't remain within the v^{th} sheet. Similarly if r is odd, we can show the action of b on (v, u_0) doesn't remain within the v^{th} sheet. Hence the alternate actions of any of (10) or (11) goes outside the sheet. □

Proposition 2.3. *If $r > 0$, $n > 1$ and $a(v) \neq v \in \Gamma_n$, then the last n alphabets of the words $a(0^r v)$ and $b(0^r v)$ are distinct from the word v .*

Proof.

$$a(0^r v) = \begin{cases} a(0^r) a(v) & \text{if } r \text{ is even} \\ a(0^r) b(v) & \text{if } r \text{ is odd} \end{cases}$$

But $a(v) \neq v \in \Gamma_n$, $\forall n > 1$ and $b(v) \neq v$, $\forall v \in \Gamma_n$, $\forall n$. Therefore the last n alphabets of the word $a(0^r v)$ are distinct from the word v . Similarly using action of b on $0^r v$ we can prove that the last n alphabets of the word $b(0^r v)$ are also distinct from the word v . $\square$

3 ZIG ZAG PRODUCT OF GRAPHS

The zig zag product of two graphs was introduced by O. Reingold, S. Vadhan and A. Wigderson[7]. Taking a product of a large graph with a small graph, the resulting graph inherits (roughly) its size from the large one, its degree from the small one, and its expansion properties from both graphs. The important property of this product is it is a good expander if both large and small graphs are expander. D. D'Angeli, A. Donno and E. Sava-Huss[3] have provided the sufficient condition for the zig-zag product of two graphs to be connected.

Definition 3.1. Let $G_1 = (V_1, E_1)$ be connected d_1 regular graph, and let $G_2 = (V_2, E_2)$ be connected d_2 regular graph such that $|V_2| = d_1$. Let Rot_{G_1} (resp. Rot_{G_2}) be the rotation map of G_1 (resp. G_2). The zig zag product $G_1 \mathbb{Z} G_2$ is a regular graph of degree d_2^2 with vertex set $V_1 \times V_2$, that we identify with the set $V_1 \times [d_1]$, and whose edges are described by the rotation map

$$\text{Rot}_{G_1 \mathbb{Z} G_2}((v, k), (i, j)) = ((w, l), (j', l')),$$

for all $v \in V_1, k \in [d_1], i, j \in [d_2]$, if :


$$(1) \text{Rot}_{G_2}(k, i) = (k', i'),$$

$$(2) \text{Rot}_{G_1}(v, k') = (w, l'),$$

$$(3) \text{Rot}_{G_2}(l', j) = (l, j'),$$

where $w \in V_1, l, k', l' \in [d_1]$ and $i', j' \in [d_2]$ and $[d_i] = \{1, 2, \dots, d_i\}$, where $i = 1, 2$

D. D'angeli, A. Donno and E. Seva-Huss have shown in Proposition 6.1 of [3] that the graph $\Gamma_n \mathbb{Z} C_4$ is isomorphic to the double cycle graph DC_{2n+1} . From the construction of the Schreier graph Γ_n of Basilica group see Figure 1, we get the natural labeling of Γ_n , we label the graph C_4 as follows:



Proposition 3.1. *Let $\Gamma_n, n \geq 1$ be the Schreier graph of the action of the Basilica group on $\{0, 1\}^n$ and C_4 be the 4 length cycle graph. Let $r \geq 0$. Then the graph $\Gamma_{n+r}(\mathbb{Z})C_4$ is an unramified graph covering of $\Gamma_r(\mathbb{Z})C_4$.*

Proof. In the Proposition 2.1, we have shown that the map $\pi : \Gamma_{n+r} \rightarrow \Gamma_r$ is a covering map. Define the map $\Pi : \Gamma_{n+r}(\mathbb{Z})C_4 \rightarrow \Gamma_r(\mathbb{Z})C_4$ as

$$\Pi(uv, s) = (\pi(uv), s) = (u, s), uv \in V(\Gamma_{n+r}), s \in \{a^{\pm 1}, b^{\pm 1}\}$$

We now show that Π is adjacency preserving map.

As,

$$\mathbf{N}_{\Gamma_{n+r}(\mathbb{Z})C_4}(uv, a) = \{((uv)^{a^{-1}}, a^{-1}), ((uv)^{a^{-1}}, b^{-1}), ((uv)^{b^{-1}}, a^{-1}), ((uv)^{b^{-1}}, b^{-1})\}$$

So we consider (uv, a) is adjacent to $((uv)^{b^{-1}}, b^{-1})$ and $\Pi(uv, a) = (u, a)$,
 $\Pi((uv)^{b^{-1}}, b^{-1}) = \Pi((u)^{b^{-1}}v', b^{-1}) = ((u)^{b^{-1}}, b^{-1}) \in \mathbf{N}_{\Gamma_r(\mathbb{Z})C_4}(u, a)$

Hence, $\Pi(uv, a)$ is adjacent to $\Pi((uv)^{b^{-1}}, b^{-1})$.

Using a similar argument we can see that other choices of s also gives that Π is adjacency preserving map.

Given $(u, s) \in V(\Gamma_r(\mathbb{Z})C_4)$, any vertex $w \in \Pi^{-1}(u, s)$ is of the form $w = (uv, s)$, for some $v \in V(\Gamma_n)$. By definition of the zig zag product of graphs, we see that the vertices in $V(\Gamma_r(\mathbb{Z})C_4)$ adjacent to (u, s) if $s = a$, have the form

$$\mathbf{N}_{\Gamma_r(\mathbb{Z})C_4}(u, a) = \{(u^{a^{-1}}, a^{-1}), (u^{a^{-1}}, b^{-1}), (u^{b^{-1}}, a^{-1}), (u^{b^{-1}}, b^{-1})\}$$

and the vertices in $V(\Gamma_{n+r}(\mathbb{Z})C_4)$ adjacent to $w = (uv, s)$ if $s = a$, have the form

$$\mathbf{N}_{\Gamma_{n+r}(\mathbb{Z})C_4}(uv, a) = \{((uv)^{a^{-1}}, a^{-1}), ((uv)^{a^{-1}}, b^{-1}), ((uv)^{b^{-1}}, a^{-1}), ((uv)^{b^{-1}}, b^{-1})\}$$

Notice that, any $s \in \{a^{\pm 1}, b^{\pm 1}\}$, $(uv)^s = s(uv)$. The length of the word $s(uv)$ is $n + r$. By the action of the s on uv , the word containing first r alphabets of $s(uv)$ is actually the word $s(u)$ and the remaining n alphabets of $s(uv)$ form some word say v' . Thus $\pi((uv)^s) = \pi(s(uv)) = \pi(s(u)v') = s(u) = u^s$.

$$\implies \Pi(\mathbf{N}_{\Gamma_{n+r}(\mathbb{Z})C_4}(uv, a)) = \mathbf{N}_{\Gamma_r(\mathbb{Z})C_4}(u, a)$$

Hence, Π is sending neighborhoods of $\Gamma_{n+r}(\mathbb{Z})C_4$ one to one and onto neighborhoods of $\Gamma_r(\mathbb{Z})C_4$ if $s = a$. Similar argument one can make in the case of $s = b, b^{-1}$ and a^{-1} . $\square$

Notice that if $Y = \Gamma_r(\mathbb{Z})C_4$, and $Y' = \Gamma_{1+r}(\mathbb{Z})C_4$,

$$\mathbf{N}_G(w, a) = \mathbf{N}_G(w, b), \mathbf{N}_G(w, a^{-1}) = \mathbf{N}_G(w, b^{-1}), \text{ for } G = Y, Y' \quad (20)$$

Let T be the spanning subgraph of Y_1 , where $V(T) = V(Y)$ and

$$E(T) = E(Y) \setminus \{e_i | i = 1, 2, 3, 4\}$$

where $u_0 = 0^r$, and, if r is even, then

$$e_1 = \{(u_0, a^{-1}), (a(b^{-1}a)^{2^r-1}u_0, a)\}, e_2 = \{(u_0, a^{-1}), (a(b^{-1}a)^{2^r-1}u_0, b)\}$$

$$e_3 = \{(u_0, b^{-1}), (a(b^{-1}a)^{2^r-1}u_0, a)\}, e_4 = \{(u_0, b^{-1}), (a(b^{-1}a)^{2^r-1}u_0, b)\}$$

and, if r is odd, then

$$e_1 = \{(u_0, a^{-1}), (b(a^{-1}b)^{2^r-1}u_0, a)\}, e_2 = \{(u_0, a^{-1}), (b(a^{-1}b)^{2^r-1}u_0, b)\}$$

$$e_3 = \{(u_0, b^{-1}), (b(a^{-1}b)^{2^r-1}u_0, a)\}, e_4 = \{(u_0, b^{-1}), (b(a^{-1}b)^{2^r-1}u_0, b)\}$$

Corollary 3.1. $\Gamma_{1+r} \otimes C_4$ is the two fold normal covering of the graph $\Gamma_r \otimes C_4$ with Galois group

$$\text{Gal}(\Gamma_{1+r} \otimes C_4 / \Gamma_r \otimes C_4) \cong \frac{\mathbb{Z}}{2\mathbb{Z}}.$$

Proof. The adjacency matrices of graphs $\Gamma_k \otimes C_4$, $k = 1 + r$, r are the circulant matrices, see [3]. Therefore there exists the following order of the vertices of $\Gamma_{1+r} \otimes C_4$ if r is even:

$$\begin{aligned} &(0^{1+r}, a^{-1}); (a(0^{1+r}), a); (b^{-1}a(0^{1+r}), a^{-1}); (ab^{-1}a(0^{1+r}), a); \\ &(b^{-1}ab^{-1}a(0^{1+r}), a^{-1}); \dots (a(b^{-1}a)^{2^{1+r}-1}(0^{1+r}), a); \end{aligned} \quad (21)$$

$$\begin{aligned} &(0^{1+r}, b^{-1}); (a(0^{1+r}), b); (b^{-1}a(0^{1+r}), b^{-1}); (ab^{-1}a(0^{1+r}), b); \\ &(b^{-1}ab^{-1}a(0^{1+r}), b^{-1}); \dots (a(b^{-1}a)^{2^{1+r}-1}(0^{1+r}), b). \end{aligned} \quad (22)$$

In other words we are applying (10) i.e. alternate actions of a and b^{-1} on the word 0^{1+r} , with the alternation of a^{-1}, a in the second coordinate of the 2^{1+r} vertices (the inner cycle of the graph) given in equation (21).

Also we are applying same action given in (10) i.e. alternate actions of a and b^{-1} on the word 0^{1+r} , with the alternation of b^{-1}, b in the second coordinate of 2^{1+r} vertices (the outer cycle of the graph) given in equation (22).

In the Proposition 2.1, we have also seen that $\Gamma_n \otimes \Gamma_r \simeq \Gamma_{n+r}$ the isomorphism is $f : \Gamma_n \otimes \Gamma_r \rightarrow \Gamma_{n+r}$ defined by $f(v, u) = uv$ and the isomorphism $f^{-1} : \Gamma_{n+r} \rightarrow \Gamma_n \otimes \Gamma_r$ defined by $f^{-1}(uv) = (v, u)$, hence we are able to write $f(0, 0^r) = 0^{1+r}$ or $f^{-1}(0^{1+r}) = (0, 0^r)$

Thus the vertex $(0^{1+r}, a^{-1})$ can be written as $((0, 0^r), a^{-1})$. But the first coordinate of this vertex i.e. $(0, 0^r)$ is the 0^r vertex of the 0^{th} sheet in the graph $\Gamma_1 \otimes \Gamma_r$, and $a(0) = 0 \in \Gamma_1$. By even case of above Proposition 2.2, we get

$$(0, 0^r), (0, a(0^r)), (0, b^{-1}a(0^r)), \dots (0, a(b^{-1}a)^{2^r-1}(0^r)) = (0, b(0^r))$$

produces spanning path of the 0^{th} sheet. But again by Proposition 2.1, and π is onto map, we have

$$\begin{aligned} f(0, 0^r) &= 0^{1+r} \\ f(0, a(0^r)) &= a(0^{1+r}) \\ f(0, b^{-1}a(0^r)) &= b^{-1}a(0^{1+r}) \\ \dots f(0, a(b^{-1}a)^{2^r-1}(0^r)) &= f(0, b(0^r)) = a(b^{-1}a)^{2^r-1}(0^{1+r}) = b(0^{1+r}) \end{aligned}$$

This gives that the first coordinate of the first 2^r vertices of the lists given in equations (21) and (22) contains 0 in the last position i.e. they are vertices of the 0^{th} sheet of $\Gamma_1 \otimes \Gamma_r$. Therefore by definition of zigzag product of graphs, equation (20) and the Proposition 2.2, these vertices forms a (connected) copy of the spanning subgraph T of Y . As the first coordinate of these vertices ends with 0, we denote this copy by T_0 . Using similar argument, we can show that first coordinate the next 2^r vertices given in (21) and (22) ends with 1 and hence we get the another copy of T which we denote by T_1 . Following are the vertices of T_0 and T_1 .

$$\begin{aligned} V(T_0) &= \{(0^{1+r}, a^{-1}); (a(0^{1+r}), a); (b^{-1}a(0^{1+r}), a^{-1}); (ab^{-1}a(0^{1+r}), a); \dots \\ &\quad (a(b^{-1}a)^{2^r-1}(0^{1+r}), a); \\ &\quad (0^{1+r}, b^{-1}); (a(0^{1+r}), b); (b^{-1}a(0^{1+r}), b^{-1}); \dots (a(b^{-1}a)^{2^r-1}(0^{1+r}), b)\} \\ &= \{(u0, s) | u \in V(\Gamma_r), s \in V(C_4)\} \end{aligned}$$

$$\begin{aligned} V(T_1) &= \{((b^{-1}a)^{2^r}0^{1+r}, a^{-1}); (a(b^{-1}a)^{2^r}(0^{1+r}), a); ((b^{-1}a)^{2^r+1}(0^{1+r}), a^{-1}); \dots \\ &\quad (a(b^{-1}a)^{2^r+1-1}(0^{1+r}), a); \\ &\quad ((b^{-1}a)^{2^r}0^{1+r}, b^{-1}); (a(b^{-1}a)^{2^r}(0^{1+r}), b); ((b^{-1}a)^{2^r+1}(0^{1+r}), b^{-1}); \\ &\quad \dots (a(b^{-1}a)^{2^r+1-1}(0^{1+r}), b)\} \\ &= \{(u1, s) | u \in V(\Gamma_r), s \in V(C_4)\} \end{aligned}$$

and

$$V(\tilde{Y}_1) = V(T_0) \cup V(T_1).$$

Define the non identity Frobenius automorphism map Σ as

$$\Sigma(u1, s) = (\sigma(u1), s) = (u0, s),$$

and

$$\Sigma(u0, s) = (\sigma(u0), s) = (u1, s), \forall u \in V(\Gamma_r), s \in V(C_4),$$

where $\sigma \in G(\Gamma_{1+r}|\Gamma_r)$.

$$\Sigma(T_0) = T_1, \Sigma(T_1) = T_0$$

and since $\sigma^2 = id$ we have

$$\Sigma^2 = Id.$$

$$(\Pi \circ \Sigma)(u0, s) = \Pi(\Sigma(u0, s)) = \Pi(\sigma(u0), s) = \Pi(u1, s) = (\pi(u1), s) = (u, s)$$

$$(\Pi \circ \Sigma)(u1, s) = \Pi(\Sigma(u1, s)) = \Pi(\sigma(u1), s) = \Pi(u0, s) = (\pi(u0), s) = (u, s)$$

and

$$\Pi(u0, s) = (u, s) = \Pi(u1, s),$$

$$\implies \Pi \circ \Sigma = \Pi.$$

Thus Galois group $G(Y'|Y) = G(\Gamma_{1+r} \otimes C_4 | \Gamma_r \otimes C_4)$ contains two elements Σ and $\Sigma^2 = Id$. One can make a similar argument for the odd n . $\square$

Corollary 3.2. *If $n > 1$, then $\Gamma_{n+r} \otimes C_4$ is the non normal covering of the graph $\Gamma_r \otimes C_4$.*

Proof of this corollary is given at the end of this paper. To show that $\Gamma_{n+r} \otimes C_4$ is not a Galois covering of $\Gamma_r \otimes C_4$ we need the following.

By definition of zigzag product of graphs,

$$V(\Gamma_{n+r} \otimes C_4) = \{(uv, s) | uv \in V(\Gamma_{n+r}), s \in V(C_4)\}$$

Let $v_0 = 0^n, v_1, v_2, \dots, v_{2^n-1}$ be the vertices of the graph Γ_n .

By Proposition 2.1, Γ_{n+r} is 2^n sheeted cover of Γ_r , so we can have following partition,

$$V(\Gamma_{n+r}) = \bigcup_{j=0}^{2^n-1} \{uv_j | u \in V(\Gamma_r)\}$$

Call

$$S_{v_j} = \{(uv_j, s) | u \in V(\Gamma_r), s \in V(C_4)\}$$

Therefore we can have

$$V(\Gamma_{n+r} \otimes C_4) = \bigcup_{j=0}^{2^n-1} S_{v_j}$$

3.1 Restriction Graph

Definition 3.2. Let $G = (V, E)$ be a graph and $\emptyset \neq V' \subsetneq V$, we define restriction graph $G' = G|_{V'}$ which is a subgraph of G as $G' = (V', E')$

$$E' = \{e = \{v_l, v_m\} \in E | v_l, v_m \in V'\}.$$

Suppose

$$\tilde{Y} = \Gamma_{n+r}(\mathbb{Z})C_4, Y = \Gamma_r(\mathbb{Z})C_4$$

we find the restriction subgraphs $\tilde{Y}_j$ of $\tilde{Y}$ by taking $V(\tilde{Y}_j) = S_{v_j}$

One can see,

$$V(\tilde{Y}) = \bigcup_{j=0}^{2^n-1} V(\tilde{Y}_j)$$

The next result is about connectedness of restriction subgraphs $\tilde{Y}_j$'s of $\tilde{Y}$.

Proposition 3.2. If $\tilde{Y}_j$'s and $\tilde{Y}$ as above, then following holds

1. If $a(v_j) = v_j \in \Gamma_n$ then the restriction subgraph $\tilde{Y}_j$ of $\tilde{Y}$ is the connected v_j^{th} sheet in the graph $\tilde{Y}$.
2. If $a(v_j) \neq v_j \in \Gamma_n$ then the restriction subgraph $\tilde{Y}_j$ of $\tilde{Y}$ is the v_j^{th} sheet in the graph $\tilde{Y}$ and this sheet is disconnected.

Proof. Recall that T is a spanning subgraph of Y , if r is even, we can write

$$V(T) = \{(0^r, a^{-1}); (a0^r, a); (b^{-1}a0^r, a^{-1}); \dots; (a(b^{-1}a)^{2^r-1}0^r, a); \\ (0^r, b^{-1}); (a0^r, b); (b^{-1}a0^r, b^{-1}); \dots; (a(b^{-1}a)^{2^r-1}0^r, b)\}$$

1. We write set T_j , $\forall v_j \in \Gamma_n$ as

$$T_j = \{(0^r v_j, a^{-1}); ((a(0^r))v_j, a); ((b^{-1}a(0^r))v_j, a^{-1}); \dots; ((a(b^{-1}a)^{2^r-1}(0^r))v_j, a); \\ (0^r v_j, b^{-1}); ((a(0^r))v_j, b); ((b^{-1}a(0^r))v_j, b^{-1}); \dots; ((a(b^{-1}a)^{2^r-1}(0^r))v_j, b)\}$$

To show that $\tilde{Y}_j$ is connected it is enough to show that $V(\tilde{Y}_j) = T_j$.

As $(0^r v_j, a^{-1}) = (u_0 v_j, a^{-1}) \in S_{v_j} = V(\tilde{Y}_j)$ and action of $b^{-1}a$ on $0^r \in X^r$ has order 2^r ,

$$\implies T_j \subset V(\tilde{Y}_j), \forall j$$

But both the sets T_j and $V(\tilde{Y}_j)$ contain 2^r elements and $T_j \subset V(\tilde{Y}_j)$.

Therefore

$$T_j = V(\tilde{Y}_j), \forall j$$

by Proposition 2.1, and equation (20) elements in T_j forms a copy of the spanning subgraph T of Y in $\tilde{Y}$, and therefore $\tilde{Y}_j$ is the connected v_j^{th} sheet in the graph $\tilde{Y}$.

2. By Proposition 2.3, $a(0^r v_j) = a(0^r) a(v_j) \neq a(0^r) v_j$ and $b(0^r v_j) \neq b(0^r) v_j$. and

$$\mathbf{N}_{\tilde{Y}}(0^r v_j, a^{-1}) = \{(a(0^r v_j), a), (a(0^r v_j), b), (b(0^r v_j), a), (b(0^r v_j), b)\}$$

Therefore

$$\mathbf{N}_{\tilde{Y}}(0^r v_j, a^{-1}) \cap \tilde{Y}_j = \emptyset,$$

which means Y_j has more than one components in $\tilde{Y}$, so $\tilde{Y}_j$ is the v_j^{th} sheet in the graph $\tilde{Y}$ and this sheet is disconnected.

□

Now we can prove the Corollary 3.2

Proof. By contradiction

Let $\tilde{Y}$ be 2^n sheeted normal covering of Y . Therefore there exists a Frobenius automorphism $\Sigma \in G(\tilde{Y}|Y)$ such that Σ sends v_j^{th} sheet in $\tilde{Y}$ to some say v_k^{th} sheet in $\tilde{Y}$, where $a(v_j) = v_j \in \Gamma_n$ and $a(v_k) \neq v_k \in \Gamma_n$.

But by above Proposition 2.4 v_j^{th} sheet is connected and under a Frobenius automorphism it is mapping to disconnected v_k^{th} sheet which is a contradiction. □

Table 6: Vertex labeling for $\Gamma_1 \mathbb{Z} C_4$

label	1	2	3	4	5	6	7	8
vertex	$(0, a^{-1})$	$(1, a)$	$(1, a^{-1})$	$(0, a)$	$(0, b^{-1})$	$(1, b)$	$(1, b^{-1})$	$(0, b)$

Table 7: Notations for sheets

Vertex set	Group element
$\{1, 2, \dots, 8\}$	$\Sigma^2 = Id$
$\{1', 2', \dots, 8'\}$	Σ

Example 3.1. The graph $\Gamma_2 \mathbb{Z} C_4$ is the normal 2 sheeted cyclic covering of the graph $\Gamma_1 \mathbb{Z} C_4$ has shown in Figure 5. We label the vertices using the Table 6. We obtain spanning subgraph of the $\Gamma_1 \mathbb{Z} C_4$ by cutting edges $e_1 = \{1, 4\}$, $e_2 = \{1, 8\}$, $e_3 = \{5, 4\}$, $e_4 = \{5, 8\}$. This gives the dashed line graph T as shown in Figure 6 In the graph $\tilde{Y}$ there are 16 vertices of which 8 ends with the alphabet 0 and remaining 8 ends with 1. By Proposition 2.1, these vertices forms connected





Figure 5: The graph $\Gamma_2 \otimes C_4$ is normal 2 sheeted cyclic covering of the graph $\Gamma_1 \otimes C_4$. The 2 sheets of $\tilde{Y} = \Gamma_2 \otimes C_4$ are copies of the spanning subgraph T of $Y = \Gamma_1 \otimes C_4$ (dashed lines). The continuous lines shown in the graph $\tilde{Y}$ are lifts of the edges e_1, e_2, e_3 and e_4 , where $e_1 = \{1, 4\}, e_2 = \{1, 8\}, e_3 = \{5, 4\}$ and $e_4 = \{5, 8\}$.

Table 8: Connections between 1-sheet and Σ -sheet in $\tilde{Y}$

Vertex	adjacent vertices in $\tilde{Y}$
1	2, 6, 4', 8'
2	1, 3, 5, 7
3	2, 4, 6, 8
4	3, 7, 1', 5'
5	2, 6, 4', 8'
6	1, 3, 5, 7
7	2, 4, 6, 8
8	3, 7, 1', 5'

sheets which we label in Table 7.

The representations of the cyclic Galois group $\{1, \Sigma\}$ are the trivial representation ρ_0 and the representation defined by $\rho(1) = 1, \rho(\Sigma) = -1$. So $Q_\rho = 2I_8$. Therefore there are two cases.

Case 1 The trivial representation $\rho(1) = 1$

We use the information given in Table 8 to write the $A(1)$ and $A(\Sigma)$, hence we have

$$A(1) = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, A(\Sigma) = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

But $A_1 = A(1) + A(\Sigma) = A_{\tilde{Y}}$ (A_1 is the adjacency matrix A_Y of Y .)

Case 2 The representation ρ

Here we find

$$A_\Sigma = A(1) - A(\Sigma)$$

$$A_{\Sigma} = \begin{pmatrix} 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ -1 & 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ -1 & 0 & 1 & 0 & -1 & 0 & 1 & 0 \end{pmatrix}$$

The eigenvalues of A_1 and A_{Σ} are given in Table 9.

Table 9: Eigenvalues of A_1 and A_{Σ}

Matrix	Eigenvalues
A_1	$4, -4, 0, 0, 0, 0, 0, 0$
A_{Σ}	$-2\sqrt{2}, -2\sqrt{2}, 2\sqrt{2}, 2\sqrt{2}, 0, 0, 0, 0$

So we can write the spectrum of $\tilde{Y} = \Gamma_2 \mathbb{Z} C_4$. See Table 10.

Table 10: spectrum of $\tilde{Y} = \Gamma_2 \mathbb{Z} C_4$

Eigenvalue	Multiplicity
± 4	1
0	10
$\pm 2\sqrt{2}$	2

Reciprocals of L functions for $\Gamma_2 \mathbb{Z} C_4 | \Gamma_1 \mathbb{Z} C_4$.

1) For A_1

$$\zeta_{\Gamma_1 \mathbb{Z} C_4}(t)^{-1} = L(t, A_1, \tilde{Y}|Y)^{-1} = (1-t^2)^8 (t-1)(t+1)(3t-1)(3t+1) (3t^2 + 1)^6.$$

2) For A_{Σ}

$$L(t, A_{\Sigma}, \tilde{Y}|Y)^{-1} = (1-t^2)^8 (3t^2 + 1)^4 (9t^4 - 2t^2 + 1)^2$$

By equation (9) we have,

$$\begin{aligned} \zeta_{\Gamma_2 \mathbb{Z} C_4}(t)^{-1} &= L(t, A_1, \tilde{Y}|Y)^{-1} L(t, A_{\Sigma}, \tilde{Y}|Y)^{-1} \\ &= (1-t^2)^{16} (t-1)(t+1)(3t-1)(3t+1) (3t^2 + 1)^6 (3t^2 + 1)^4 (9t^4 - 2t^2 + 1)^2 \end{aligned}$$

Table 11: Notations of sheets

Vertex set	Sheet index
$\{1, 2, \dots, 8\}$	1^{st}
$\{1', 2', \dots, 8'\}$	2^{nd}
$\{1'', 2'', \dots, 8''\}$	3^{rd}
$\{1''', 2''', \dots, 8'''\}$	4^{th}

Example 3.2. Figure 6 contains the example of non normal covering.

We obtain spanning subgraph of the $\Gamma_1(\mathbb{Z})C_4$ by cutting edges $e_1 = \{1, 4\}, e_2 = \{1, 8\}, e_3 = \{5, 4\}, e_4 = \{5, 8\}$. This gives the dashed line graph T as shown in Figure 6. In the graph $\tilde{Y}$ there are 16 vertices of which 8 ends with the alphabet 0 and remaining 8 ends with 1. The labeling of vertices is given in Table 11. By Proposition 2.1, these vertices forms connected sheets. The labeling of sheets is given in Table 12.

Table 12: Connections between 2^{nd} -sheet in $\tilde{Y}$ and other sheets

Vertex	adjacent vertices in $\tilde{Y}$
$1'$	$4, 8, 4'', 8''$
$2'$	$1'', 3', 5'', 7'$
$3'$	$2', 4', 6', 8'$
$4'$	$3', 7', 1, 5$
$5'$	$4, 8, 4'', 8''$
$6'$	$1'', 3', 5'', 7'$
$7'$	$2', 4', 6', 8'$
$8'$	$3', 7', 1, 5$

Table 13: Vertex labeling for $\Gamma_1(\mathbb{Z})C_4$

label	1	2	3	4	5	6	7	8
vertex	$(0, a^{-1})$	$(1, a)$	$(1, a^{-1})$	$(0, a)$	$(0, b^{-1})$	$(1, b)$	$(1, b^{-1})$	$(0, b)$

From the Table 13 we can easily observe the following,

$$N_{\tilde{Y}}(1') = \{4, 8, 4'', 8''\} \cap \tilde{Y}_2 = \emptyset$$




Figure 6: The graph $\Gamma_3 \otimes C_4$ is a non normal 4 sheeted covering of the graph $\Gamma_1 \otimes C_4$. The 4 sheets of $\Gamma_3 \otimes C_4$ are copies of the spanning subgraph T of $\Gamma_1 \otimes C_4$ (dashed lines).

By Corollary 3.2, the graph $\Gamma_3 \mathbb{Z} C_4$ is non normal 4 sheeted covering of the graph $\Gamma_1 \mathbb{Z} C_4$.

We propose the following conjecture on the basis of this study. The work presented in this paper may be the special case of this conjecture.

Conjecture 3.1. *If $\tilde{G}$ is normal covering of G with Galois group $Gal(\tilde{G}|G)$ and H be some other graph, then $\tilde{G} \mathbb{Z} H$ is also normal covering of $G \mathbb{Z} H$ with Galois group $Gal(\tilde{G} \mathbb{Z} H|G \mathbb{Z} H) \simeq Gal(\tilde{G}|G)$.*

References

- [1] A. Abdollahi, & and A. Loghman, On one-factorizations of replacement products, *Filomat*, 27(1),(2013), 57-63.
- [2] D. D'Angeli, A. Donno, M. Matter and T. Nagnibeda, Schreier graphs of the basilica group, *arXiv preprint arXiv:0911.2915* (2009), *Journal of Modern Dynamics*, 4(1), (2010), 139-177.
- [3] D. D'Angeli, A. Donno and E. Sava-Huss, Connectedness and isomorphism properties of the zig-zag product of graphs, *arXiv preprint arXiv:1404.4342* (2014).
- [4] M. DeDeo, . Martínez, A. Medrano, M. Minei, H. Stark and A. Terras Michelle, et al, *Zeta Functions of Heisenberg Graphs over Finite Rings, Theory and Applications of Special Functions* Mourad E. H. Ismail, Erik Koelink,(Editors),Springer(2005), 165-183.
- [5] R. I. Grigorchuk,, and A. Żuk, On a torsion-free weakly branch group defined by a three state automaton, *International Journal of Algebra and Computation*, 12(01n02),(2002), 223-246.
- [6] R. I. Grigorchuk, Some topics in the dynamics of group actions on rooted trees, *Proceedings of the Steklov Institute of Mathematics* 273.1 (2011), 64-175.
- [7] O. Reingold, and S. Vadhan, & A. Wigderson, Entropy waves, the zig-zag graph product, and new constant-degree expanders, *Annals of mathematics*, (2002) 157-187.
- [8] A. Terras, *Zeta functions of graphs: a stroll through the garden* (Vol. 128). Cambridge University Press, 2010.